


Y9 KEY KNOWLEDGE / LITERACY (EUCLID)

(LEARN THIS KEY INFORMATION FROM PREVIOUS YEARS)



1.	<p>A ratio is the relationship between two or more numbers that are separated by a colon</p> <p style="text-align: center;">$2 : 3$</p> <p style="text-align: center;"> </p>
2.	<p>π (or pi) is a constant that is the ratio of a circle's circumference to its diameter (approximately 3.14)</p>
3.	<p>The general form of the equation for a straight line is $y = mx + c$</p>
4.	<p>In $y = mx + c$, m represents the gradient (the steepness of the line)</p>
5.	<p>In $y = mx + c$, c represents the y-intercept (where the line crosses the y-axis)</p>
6.	<p>A negative gradient represents a line going "down-hill"</p> <div style="text-align: center;"> </div>
7.	<p>Direct proportion; as one amount increases, another amount increases at the same rate</p>
8.	<p>Direct proportion can be written in the form $y = kx$ where k is the constant of proportionality (similar to a scale factor, k is also the gradient of the line)</p>
9.	<p>A direct proportion graph is a straight line that passes through the origin (0,0)</p> <div style="text-align: center;"> </div>
10.	<p>Similar shapes in maths are enlargements of each other, their lengths are in direct proportion. (Angles remain the same in similar shapes)</p>
11.	<p>A scale factor tells us what multiplier has been used to enlarge a shape (remember enlargements can get smaller too, with scale factors between 0 and 1)</p>
12.	<p>A map scale is often written in the form 50 000:1, this means 1 cm on the map is equivalent to 50 000cm in real life.</p>
13.	<p>To multiply fractions together, multiply the numerators together and the denominators together.</p>
14.	<p>To divide fractions together, use KFC. Keep the first fraction the same, Flip the second fraction over, Change the sign to a multiply, then work it out.</p> <div style="text-align: center;"> </div>
15.	<p>A reciprocal is 1 divided by the number given, or when given as a fraction the numerator and denominator switch places.</p>






16. The **midpoint** of a line segment is the exact middle of the line.

$$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

17. We use a **scatter graph** to see if two things have a relationship.



19. **Continuous** data can take any value, often it will be a measurement.

20. **Discrete** data can only take certain values, like shoe sizes.

21. **Qualitative** data use words instead of numbers, like someone's favourite colour.

22. The **Product rule** find the total number of possible combinations of items from different groups. If one group has m items in it, and another group has n items in it, the total number of possible combinations is $m \times n$.

23. A **formula** is a rule written using symbols that represents different amounts. For example, the area of a triangle can be found with the formula $A = \frac{bh}{2}$

24. An **expression** is a collection of algebraic terms that are being added or subtracted together.

25. An **equation** shows that two things are equal.

26. An **identity** is true no matter what values are chosen. For example, $3(x + 5) \equiv 3x + 15$. \equiv is the mathematical symbol for an identity.

27. To **expand** an expression means to multiply out the brackets.

Expand the following expression

$$3(x - 9)$$

	x	-9
3	$3x$	-27

Final Ans: $3x - 27$





28. To **factorise** an expression is the reverse of expanding. So factorising $3x - 27$ gives $3(x - 9)$. For it to be fully factorised we must have the highest common factor in front of the bracket.

29. **Inequalities...**

Greater than $>$ Greater than or equal to \geq
 Less than $<$ Less than or equal to \leq
 Not equal to \neq

30.	A sequence is a set of numbers that follow a rule to get from one number to the next.
31.	The nth term is an algebraic rule that enables us to find any term in a sequence.
32.	Laws of indices, multiplying: $a^m \times a^n = a^{m+n}$
33.	Laws of indices, dividing: $a^m \div a^n = a^{m-n}$
34.	Laws of indices, powers of powers: $(a^m)^n = a^{mn}$
35.	Percentage multipliers use decimal equivalents of percentages and multiplication to calculate amounts. For example, the percentage multiplier for 63% is $\times 0.63$
36.	When calculating a percentage increase add the percent to 100 and change it to a decimal, then multiply. For example, to increase by 15% do $100+15=115$, 115% as a decimal is 1.15, so multiply the amount by 1.15
37.	When calculating a percentage decrease subtract the percent from 100 and change it to a decimal, then multiply. For example, to decrease by 15% do $100-15=85$, 85% as a decimal is 0.85, so multiply the amount by 0.85
38.	To find an original amount after a percentage change we need to find the reverse percentage . This means we divide by the multiplier . For example, to find the original amount after it was increased by 15%, we divide our answer by 1.15
39.	A number written in standard form is a number written between 1 and 10 multiplied by 10 to an appropriate power. We use standard form to represent very large or very small numbers. For example, $0.00032 = 3.2 \times 10^{-4}$, $320000 = 3.2 \times 10^5$
40.	A negative power represents the reciprocal of a number (i.e. when we flip the second fraction when dividing fractions)
41.	When dealing with a fractional power , the denominator tells us what root to take of the number, and the numerator tells us what power to take, i.e. $x^{\frac{m}{n}} = (\sqrt[n]{x})^m \text{ or } \sqrt[n]{(x^m)}$
42.	An error interval describes the range of values an answer must be between. For example, a number rounded to the nearest 10 is 100, the error interval for that number is $95 \leq x < 105$
43.	<p>Angles on parallel lines...</p> <p>Look for an F-shape Look for a Z-shape Look for a C- or U-shape</p>
44.	<p>Area of a parallelogram...</p> <p>$A = bh$</p>



45.	Area of a triangle...  $A = \frac{1}{2} b h$
46.	Area of a trapezium...  $A = \frac{(a+b)}{2} h$
47.	Area of a circle...  $A = \pi r^2$
48.	Circumference of a circle...  $C = \pi D$
49.	The range is the largest number subtract the smallest number.
50.	The mode is the <u>most</u> common number.
51.	The median is the middle number, when put in order.
52.	The mean is the total of all the numbers divided by the amount of numbers.